

Cancer Association of South Africa (CANSA)



CANSA Fact Sheet on Paget's Disease of the Breast

Introduction

The breast is the tissue overlying the chest (pectoral) muscles. Women's breasts are made of specialised tissue that produces milk (glandular tissue) as well as fatty tissue. The amount of fat determines the size of the breast.

[Picture Credit: Paget's Disease of the Breast I]



The milk-producing part of the breast is organised into 15 to 20 sections, called lobes. Within each lobe are smaller structures, called lobules, where milk is produced. The milk travels through a network of tiny tubes called ducts. The ducts connect and come together into larger ducts, which eventually exit the skin in the nipple. The dark area of skin surrounding the nipple is called the areola.

Connective tissue and ligaments provide support to the breast and give it its shape. Nerves provide sensation to the breast. The breast also contains blood vessels, lymph vessels, and lymph nodes.

[Picture Credit: Paget's Disease Male Breast]

Breast Conditions

- Breast cancer - malignant (cancer) cells multiplying abnormally in the breast, eventually spreading to the rest of the body if untreated. Breast cancer occurs almost exclusively in women, although men can be affected. Signs of breast cancer include a lump, bloody nipple discharge, or skin changes.
- Ductal carcinoma in situ (DCIS) - breast cancer in the duct cells that has not invaded deeper or spread through the body. Women diagnosed with DCIS have a high likelihood of being cured.



Researched and Authored by Prof Michael C Herbst

[D Litt et Phil (Health Studies); D N Ed; M Art et Scien; B A Cur; Dip Occupational Health; Dip Genetic Counselling; Diagnostic Radiographer; Dip Audiometry and Noise Measurement; Medical Ethicist]

Approved by Ms Elize Joubert, Chief Executive Officer [BA Social Work (cum laude); MA Social Work]

June 2025

- Lobular carcinoma in situ (LCIS) - although called a carcinoma LCIS, which occurs in the milk-producing lobule cells, does not invade or spread and is not a true cancer. However, women with LCIS have an increased likelihood of developing invasive breast cancer in the future.
- Invasive ductal carcinoma - breast cancer that begins in the duct cells but then invades deeper into the breast, carrying the potential of spreading to the rest of the body (metastasising). Invasive ductal carcinoma is the most common type of invasive breast cancer.
- Invasive lobular carcinoma - breast cancer that begins in the milk-producing lobule cells, but then invades deeper into the breast, carrying the potential of spreading to the rest of the body (metastasising). Invasive lobular carcinoma is an uncommon form of breast cancer.
- Simple breast cyst – a benign (noncancerous), fluid-filled sac that commonly develops in women in their 30s or 40s. Breast cysts may cause tenderness and may be drained.
- Breast fibroadenoma - a very common noncancerous solid tumour of the breast. A typical fibroadenoma creates a painless, mobile lump in the breast and most commonly occurs in women in their 20s or 30s.
- Fibrocystic breast disease - a common condition in which noncancerous breast lumps may become uncomfortable and change in size throughout the menstrual cycle.
- Usual hyperplasia of the breast - a breast biopsy may show normal-appearing, noncancerous ductal cells multiplying abnormally. The presence of usual hyperplasia may slightly increase a woman's lifetime risk of breast cancer.
- Atypical hyperplasia of the breast - abnormal-appearing cells multiplying either in the breast ducts (atypical ductal hyperplasia) or lobules (atypical lobular hyperplasia), sometimes discovered by a breast biopsy. Although the condition is noncancerous, women with atypical hyperplasia are at four to five times higher risk of developing breast cancer compared to women with no breast abnormalities.
- Intraductal papilloma - a noncancerous, wart-like breast mass that grows inside the breast ducts. Intraductal papillomas may be felt as a lump or cause clear or bloody fluid to leak from the nipple.
- Adenosis of the breast: A noncancerous enlargement of the breast lobules. Adenosis can look like breast cancer on mammograms, so a biopsy may be needed to rule out breast cancer.
- Phyllodes tumour - a rare, usually large, rapidly growing breast tumour that looks like a fibroadenoma on ultrasound. Phyllodes tumours may be benign or malignant and most commonly develop in women in their 40s.
- Fat necrosis - in response to an injury in the fatty part of the breast, a lump of scar tissue may develop. This mass can seem like breast cancer on examination or in mammograms.
- Mastitis - inflammation of the breast, causing redness, pain, warmth, and swelling. Nursing mothers are at higher risk for mastitis, which is usually the result of infection.
- Breast calcifications - calcium deposits in the breast are a common finding on mammograms. The pattern of calcium might suggest cancer, leading to further tests or a biopsy.
- Gynaecomastia overdevelopment of male breasts. Gynecomastia can affect newborns, boys, and men.

Paget's Disease of the Breast

Paget's disease of the breast is a rare form of breast cancer. Paget's disease of the breast starts on the nipple and extends to the dark circle of skin (areola) around the nipple. It is not related to Paget's disease of the bone, which is a metabolic bone disease.

Paget's disease of the breast occurs most often in women older than age 50. Most women with Paget's disease of the breast have underlying ductal breast cancer, either *in situ* or, less commonly, invasive breast cancer. Only in rare cases is Paget's disease of the breast confined to the nipple itself.

Paget's disease is much more frequent in women but can also occur in men.

Researched and Authored by Prof Michael C Herbst

[D Litt et Phil (Health Studies); D N Ed; M Art et Scien; B A Cur; Dip Occupational Health; Dip Genetic Counselling; Diagnostic Radiographer; Dip Audiometry and Noise Measurement; Medical Ethicist]

Approved by Ms Elize Joubert, Chief Executive Officer [BA Social Work (cum laude); MA Social Work]

June 2025

Oba, T., Maeno, K., Iji, R., Kiyosawa, N., Park, Y., Morikawa, H., Amitani, M., Shimizu, T., Ono, M., Ito, T., Kanai, T., Matoba, H., Takayama, F. & Ito, K.I. 2025.

Purpose: Contrast-enhanced magnetic resonance imaging (CE-MRI) is an essential imaging modality for planning breast cancer surgical procedures. However, CE-MRI findings in Paget's disease of the breast (PD) have not been studied extensively. This study aimed to elucidate the CE-MRI findings of PD and assess their role in surgical treatment planning.

Methods: Twelve patients with PD who underwent surgery between 2011 and 2023 were retrospectively analyzed.

Results: The average patient age was 73.8 ± 14.3 years. CE-MRI detected enhanced areas in the nipple-areola complex (NAC) and/or surrounding skin in all patients. Additionally, 6 patients showed enhanced areas within the breast, suggesting ductal spread into the breast. Of these, 1 patient underwent breast-conserving surgery (BCS), and 5 opted for mastectomy. Pathology confirmed the extent of ductal spread of PD, as indicated by CE-MRI. Among the 6 patients who did not have an enhanced area in the breast, 3 underwent BCS or central lumpectomy including NAC, and 3 chose mastectomy based on the patient's preference, and no malignant foci were observed in the breast.

Conclusion: CE-MRI effectively evaluated the ductal spread of PD in the breast, demonstrating its utility in guiding the selection of surgical procedures.

Conflict of interest statement

Declarations. Conflicts of interest: The authors declare no conflicts of interest in association with the present study.

Slama, L., Taheri, H., Miry, N., Bennani, A., Chatbi, Z., Bellajdel, I., Saadi, H. & Mimouni, A. 2025.

Introduction: Tumors of the nipple-areolar complex are relatively uncommon, and Paget's disease is an uncommon presentation involving epidermal invasion of the nipple by a carcinoma in situ. Mammary Paget's disease (MPD) presents a diagnostic challenge, especially in the absence of a palpable breast mass, which often leads to delayed diagnosis and treatment.

Case report: We present the case of a 64-year-old woman with persistent skin erosion of the left nipple that was unresponsive to multiple topical treatments. Diagnostic evaluation, including mammography and MRI, revealed no invasive malignancy. Nipple biopsy confirmed MPD with underlying in situ ductal carcinoma. The patient underwent lumpectomy with removal of the nipple-areolar complex, followed by radiotherapy, resulting in a favorable outcome.

Discussion: Diagnosis of MPD can be challenging because of its resemblance to benign dermatological conditions. Imaging, particularly mammography and MRI, is essential for assessing the underlying malignancy and surgical planning. Treatment typically involves surgical excision, with breast-conserving therapy considered for localized disease. Early detection significantly improves the prognosis, particularly in the absence of invasive cancer.

Conclusion: MPD should be considered in patients with persistent eczematous nipple lesions resistant to topical treatments. Early diagnosis and appropriate surgical management are crucial to achieve favorable outcomes.

Conflict of interest statement

The authors have no conflicts of interest to declare.

The International Classification of Disease 10th Version (ICD-10)

The International Classification of Diseases (ICD) is designed to promote international comparability in the collection, processing, classification, and presentation of mortality statistics. This includes providing a format for reporting causes of death on the death certificate.

Researched and Authored by Prof Michael C Herbst

[D Litt et Phil (Health Studies); D N Ed; M Art et Scien; B A Cur; Dip Occupational Health; Dip Genetic Counselling; Diagnostic Radiographer; Dip Audiometry and Noise Measurement; Medical Ethicist]

Approved by Ms Elize Joubert, Chief Executive Officer [BA Social Work (cum laude); MA Social Work]

June 2025

ICD serves a broad range of uses globally and provides critical knowledge on the extent, causes and consequences of human disease and death worldwide via data that is reported and coded with the ICD. Clinical terms coded with ICD are the main basis for health recording and statistics on disease in primary, secondary and tertiary care, as well as on cause of death certificates. These data and statistics support payment systems, service planning, administration of quality and safety, and health services research. Diagnostic guidance linked to categories of ICD also standardizes data collection and enables large scale research.

For more than a century, the International Classification of Diseases (ICD) has been the basis for comparable statistics on causes of mortality and morbidity between places and over time. Originating in the 19th century, the latest version of the ICD, ICD-11, was adopted by the 72nd World Health Assembly in 2019 and came into effect on 1st January 2022.

The ICD-10 Code malignant neoplasm of nipple and areola, female (Paget's disease) – C50.01.

Incidence of Paget's Disease of the Breast

The latest edition of the National Cancer Registry (2023) does not provide any information regarding the incidence of Paget's Disease of the Breast.

Causes and Risk Factors for Paget's Disease of the Breast

Doctors do not know what causes Paget's disease of the breast. The most widely accepted theory is that the disease results from an underlying ductal breast cancer. The cancer cells from the original tumour then travel through milk ducts to the nipple and its surrounding skin.

Another theory is that the disease can develop independently in the nipple.



[Picture Credit: Paget's Disease of the Breast II]

Risk factors that affect one's likelihood of developing Paget's disease of the breast are the same factors that affect one's risk of developing any other type of breast cancer.

Some factors that make you more susceptible to breast cancer include:

- **Age.** One's chances of developing breast cancer increase as one gets older.
- **A personal history of breast cancer.** If someone has had breast cancer in one breast, he/she has an increased risk of developing cancer in the other breast.
- **A personal history of breast abnormalities.** If one has had lobular carcinoma *in situ* or atypical hyperplasia, the risk of developing breast cancer is higher. Certain benign breast conditions also are associated with a slightly increased risk.

- **Family history.** If one has a mother, sister or daughter with breast or ovarian cancer or both, or even a father or brother with breast cancer, one also has a greater chance of developing breast cancer.
- **An inherited gene mutation that increases the risk of breast cancer.** Defects in one of several genes, especially BRCA1 or BRCA2, puts one at greater risk of developing breast cancer as well as ovarian and other cancers. Such defects account for fewer than 1 out of 10 breast cancers.
- **Dense breast tissue.** Women with dense breast tissue, as seen on a mammogram, face a higher risk of breast cancer.
- **Radiation exposure.** If one received radiation treatments to one's chest as a child or young adult to treat another cancer, one is more likely to develop breast cancer later in life.
- **Excess weight.** Weighing more than is healthy for age and height increases the risk of breast cancer - especially after menopause and if one has gained weight as an adult.
- **Hormone replacement.** Taking oestrogen after menopause increases the risk of breast cancer for some women.
- **Race.** White women are more likely to develop breast cancer than black or Hispanic women, but black women are more likely to die of the disease.
- **Alcohol.** Alcohol consumption increases the risk of developing breast cancer.

Having one or more risk factors does not necessarily mean one will develop breast cancer. Most women with breast cancer have no known risk factors.

Symptoms and Diagnosis of Paget's Disease of the Breast

Paget disease of the breast (also known as Paget disease of the nipple and mammary Paget disease) is a rare type of cancer involving the skin of the nipple and, usually, the darker circle of skin around it, which is called the areola. Paget's disease is much more frequent in women but can occur in men.

Paget's disease causes the skin on and around the nipple to become red, sore, and flaky, or scaly. At first, these symptoms tend to come and go.

Over time, symptoms of Paget's disease usually worsen and may include:

- itching, tingling, and/or a burning sensation
- pain and sensitivity
- scaling and thickening of the skin
- flattening of the nipple
- yellowish or bloody discharge from the nipple

Because Paget's disease of the nipple is rare, doctors often mistake it for eczema (severe skin rash and inflammation), an infection or injury, or some other skin condition. For many people, it can take several months to get a correct diagnosis. If you have any of the above symptoms and they persist in spite of treatment, get them checked out by a breast specialist. In most cases, Paget's disease affects one breast, not both.

Diagnosis and Treatment of Paget's Disease of the Breast

Diagnosing Paget's disease usually involves the following steps:

Researched and Authored by Prof Michael C Herbst

[D Litt et Phil (Health Studies); D N Ed; M Art et Scien; B A Cur; Dip Occupational Health; Dip Genetic Counselling; Diagnostic Radiographer; Dip Audiometry and Noise Measurement; Medical Ethicist]

Approved by Ms Elize Joubert, Chief Executive Officer [BA Social Work (cum laude); MA Social Work]

June 2025

- A physical examination of the breasts, with special attention paid to the area around the nipple. The doctor may be able to feel a lump or mass in the breast.
- A mammogram to check the nipple area and also to look for evidence of cancer in other areas of the breast.
- Ultrasound and/or breast MRI to create additional images of the breast and check for other areas of cancer.
- Biopsy of the nipple and areola. A breast surgeon will perform minor surgery to remove a small piece of tissue from the nipple and areola area and examine it under a microscope. If there is unusual discharge from the breast, the doctor will take a sample of that for examination as well.

For some women with Paget's disease of the nipple, radiotherapy is almost always recommended, and may be the only treatment needed after biopsy and breast conserving surgery. Radiotherapy is sometimes also recommended after mastectomy.

For many years, mastectomy, with or without the removal of lymph nodes under the arm on the same side of chest (known as axillary lymph node dissection), was regarded as the standard surgery for Paget disease of the breast. This type of surgery was done because patients with Paget disease of the breast were almost always found to have one or more tumours inside the same breast. Even if only one tumour was present, that tumour could be located several centimetres away from the nipple and areola and would not be removed by surgery on the nipple and areola alone.

Studies have shown, however, that breast-conserving surgery that includes removal of the nipple and areola, followed by whole-breast radiation therapy, is a safe option for people with Paget disease of the breast who do not have a palpable lump in their breast and whose mammograms do not reveal a tumour.

The prognosis, or outlook, for people with Paget disease of the breast depends on a variety of factors, including the following:

- Whether or not a tumour is present in the affected breast
- If one or more tumours are present in the affected breast, whether those tumours are ductal carcinoma in situ or invasive breast cancer
- If invasive breast cancer is present in the affected breast, the stage of that cancer is staged the same as breast cancer

The presence of invasive cancer in the affected breast and the spread of cancer to nearby lymph nodes are associated with reduced survival.

Medical Disclaimer

This Fact Sheet is intended to provide general information only and, as such, should not be considered as a substitute for advice, medically or otherwise, covering any specific situation. Users should seek appropriate advice before taking or refraining from taking any action in reliance on any information contained in this Fact Sheet. So far as permissible by law, the Cancer Association of South Africa (CANSA) does not accept any liability to any person (or his/her dependants/ estate/heirs) relating to the use of any information contained in this Fact Sheet.

Whilst CANSA has taken every precaution in compiling this Fact Sheet, neither it, nor any contributor(s) to this Fact Sheet can be held responsible for any action (or the lack thereof) taken by any person or organisation wherever they shall be based, as a result, direct or otherwise, of information contained in, or accessed through, this Fact Sheet.



Sources and References Consulted or Utilised

Adams, S.J. & Kanthan, R. 2016. Paget's disease of the male breast in the 21st century: A systematic review. *Breast*. 2016 Oct;29:14-23. doi: 10.1016/j.breast.2016.06.015. Epub 2016 Jul 6.

BreastCancer.Org

http://www.breastcancer.org/symptoms/types/pagets/sympt_diag

Cancer Research UK

<http://www.cancerresearchuk.org/about-cancer/breast-cancer/stages-types-grades/types/pagets-disease-breast>

Dubar, S., Boukrid, M., Bouquet de Joliniere, J., Guillou, L., Vo, Q>D>, Major, A., Alin N.B. Khomsi, F. & Feki, A. 2017. Paget's Breast Disease: A Case Report and Review of the Literature. *Front Surg*. 2017 Oct 23;4:51. doi: 10.3389/fsurg.2017.00051. eCollection 2017.

Gabrielová, L., Hanincová, V., Coufal, O., Holánek, M. & Zapletal, O. 2021. Pagets disease of the breast - observational retrospective study of the past 20 years. *Rozhl Chir*. Spring 2021;100(4):173-179.

Gaurav, A., Gupta, V., Koul, R., Dabas, S., Sareen, R., Geeta, K., Arora, V., Parikh, P.M. & Aggarwal, S. 2018. Practical consensus recommendatons for Paget's disease in breast cancer. *South Asian J Cancer*. 2018 Apr-Jun;7(2):83-86. doi: 10.4103/sajc.sajc_107_18.

Godbole C, Mehta J, Methil B, Palep R, Bhuta P. 2017. Perianal Paget's Disease-a Case Report and a Review of Current Diagnosis and Management. *Indian J Surg Oncol*. 2017 Dec;8(4):619-621. doi: 10.1007/s13193-016-0548-7. Epub 2016 Sep 1.

Grewal, I.S. & Rajagopal, T. 2023. Studying the presentation of Paget disease of the breast. *Cureus*. 2023 Jun 27;15(6):e41057.

Hudson-Phillips, S., Cox, K., Patel, P. & Al Sarakbi, W. 2023. Paget's disease of the breast: diagnosis and management. *Br J Hosp Med (Lond)*. 2023 Jan 2;84(1):1-8.

International Classification of Diseases 10th version (ICD-10)

[https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2842174/#:~:text=Malignant%20rhabdoid%20tumors%20\(MRTs\)%20are,particularly%20in%20the%20soft%20tissues](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2842174/#:~:text=Malignant%20rhabdoid%20tumors%20(MRTs)%20are,particularly%20in%20the%20soft%20tissues).

<https://www.dana-farber.org/childhood-malignant-rhabdoid-tumor/>

<https://medlineplus.gov/genetics/condition/rhabdoid-tumor-predisposition-syndrome/>

<https://www.hindawi.com/journals/crpu/2014/950869/>

<https://jmedicalcasereports.biomedcentral.com/articles/10.1186/s13256-017-1554-2>

<http://www.cancerindex.org/ccw/guide3g.htm>

<https://www.dana-farber.org/childhood-malignant-rhabdoid-tumor/#:~:text=Diagnostic%20procedures%20for%20malignant%20rhabdoid,to%20test%20for%20genetic%20mutation>.

Kenny, M.G., Nguyen, M., Modi, M., Eilers, D. & Souchik, A. 2024. Melanoma Mimicker: Pigmented Mammary Paget Disease in a Man. *Am J Dermatopathol*. 2024 Jan 1;46(1):46-49.

Li, C., Yang, X., Wang, P., We, L. & Wang, X. 2020. Mammary Paget's disease diagnosed with reflectance confocal microscopy. *Photodiagnosis Photodyn Ther*. 2020 Dec;32:102069.

Markarian, S. & Holmes, D.R. 2022. Mammary Paget's Disease: an update. *Cancers (Basel)*. 2022 May 13;14(10):2422.

Researched and Authored by Prof Michael C Herbst

[D Litt et Phil (Health Studies); D N Ed; M Art et Scien; B A Cur; Dip Occupational Health; Dip Genetic Counselling; Diagnostic Radiographer; Dip Audiometry and Noise Measurement; Medical Ethicist]

Approved by Ms Elize Joubert, Chief Executive Officer [BA Social Work (cum laude); MA Social Work]

June 2025

Mayo Clinic

<https://www.mayoclinic.org/diseases-conditions/pagets-disease-of-the-breast/symptoms-causes/syc-20351079>

Moore, S.A., Notgrass, H.M., Vandergriff, T.W. & Sahoo, S. 2020. Mammary Paget's disease of the male breast: a rare case with an unusual immunohistochemical profile. *Int J Surg Pathol.* 2020 Apr;28(2):210-215. doi: 10.1177/1066896919874878. Epub 2019 Sep 12.

Mousa-Doust, D., Warburton, R., Pao, J.S., Dingee, C., Bazzarelli, A., Newman-Bremang, J. & McKeivitt, E. 2024. Paget's disease of the breast: Presentation, treatment, and outcomes in a modern cohort. *Am J Surg.* 2024 May;231:18-23.

National Cancer Institute

<https://www.cancer.gov/types/breast/paget-breast-fact-sheet#q6>

Nori, J., Bicchierai, G., Amato, F., De Benedetto, D., Boeri, C., Vanzi, E., Di Naro, F., Bianchi, S. & Miele, V. 2021. A new technique for the histological diagnosis of Paget's disease of the breast using a semiautomated core needle biopsy with a 14-gauge needle. *Radiol Med.* 2021 Jul;126(7):936-945.

Oba, T., Maeno, K., Iji, R., Kiyosawa, N., Park, Y., Morikawa, H., Amitani, M., Shimizu, T., Ono, M., Ito, T., Kanai, T., Matoba, H., Takayama, F. & Ito, K.I. 2025. Utility of contrast-enhanced magnetic resonance imaging for planning of surgical procedure in Paget's disease of the breast. *Surg Today.* 2025 Jun;55(6):778-786.

Ooi, P.S., Draman, N., Yusoff, S.S.M., Zain, W.Z.W., Ganasagaran, D. & Chua, H.H. 2018. Mammary Paget's Disease of the Nipple: Relatively Common but Still Unknown to Many. *Korean J Fam Med.* 2018 Nov 29. doi: 10.4082/kjfm.17.0143. [Epub ahead of print]

Owusu-Brackets, N., Chen, J.C., Li, Y., Fisher, J.L., Bhattacharyya, O. & Obeng-Gyasi, S. 2022. Examining racial differences in treatment and survival among patients with Paget's disease of the breast. *Surgery.* 2022 Oct 20;S0039-6060(22)00713-9.

Paget's Disease Male Breast

<http://www.bioline.org.br/pdf?cr10021>

Paget's Disease of the Breast I

<http://www.pathologyoutlines.com/topic/breastmalignantpaget.html>

Paget's Disease of the Breast II

<https://radiopaedia.org/articles/paget-disease-of-the-breast-1>

Pourmousa, A.J., Mautner, S.K., Nasseri-Nik, N. & Lampen-Sachar, K. 2021. Invasive Paget's Disease of the Breast: Rash or Recurrence? *Cureus.* 2021 Nov 2;13(11):e19216.

Radiopaedia

<https://radiopaedia.org/articles/paget-disease-of-the-breast-1>

RSNA RadioGraphics

<http://pubs.rsna.org/doi/full/10.1148/rg.317115070>

Salih, A.M., Hammood, Z.D., Kakamad, F.H., Othman, S., Ali, R.K. & Latif, S. 2021. Paget's disease of the breast in male with underlying invasive ductal carcinoma: A case report with review of literature. *Ann Med Surg (Lond).* 2021 Nov 11;72:103035.

Slama, L., Taheri, H., Miry, N., Bennani, A., Chatbi, Z., Bellajdel, I., Saadi, H. & Mimouni, A. 2025. Mammary Paget's Disease of the Nipple: A Case Report. *Case Rep Oncol.* 2025 May 26;18(1):780-785.

Uthamalingam, M. & Periyasamy, K. 2016. Paget's Disease of Nipple in Male Breast with Cancer. *J Clin Diagn Res.* 2016 Feb; 10(2): PD14-PD16.

WebMD

<https://www.webmd.com/women/picture-of-the-breasts#1>

Yao, Y., Sun, L., Meng, Y., Zhuang, Y., Zhao, L., Yu, Q. & Si, C. 2019. Breast-conserving surgery in patients with mammary Paget's disease. *J Surg Res.* 2019 Sep;241:178-187. doi: 10.1016/j.jss.2019.03.025. Epub 2019 Apr 24.

Researched and Authored by Prof Michael C Herbst

[D Litt et Phil (Health Studies); D N Ed; M Art et Scien; B A Cur; Dip Occupational Health; Dip Genetic Counselling; Diagnostic Radiographer; Dip Audiometry and Noise Measurement; Medical Ethicist]

Approved by Ms Elize Joubert, Chief Executive Officer [BA Social Work (cum laude); MA Social Work]

June 2025

Yasir, M., Khan, M. & Lotfollahzadeh, S. 2022. Mammary Paget Disease. *In*: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan. 2022 Sep 26.